



ARBOR ASSAYS
Interactive Assay Solutions™

STRESS ASSAY KITS

WEB 180403

Table of Contents

Assay Kits	Page #
Catalase Colorimetric and Fluorescent Activity Kits	3
Ceruloplasmin Colorimetric Activity Kit	4
Corticosterone EIA and CLIA Kits	5
Cortisol EIA Kits	6
Cortisone EIA and CLIA Kits	7
FRAP™ (Ferric Reducing Antioxidant Power) Detection Kit	8
Glutathione Colorimetric Detection Kit	9
Glutathione Fluorescent Detection Kits	10
Glutathione Reductase Fluorescent Activity Kit	11
Glutathione S-Transferase (GST) Fluorescent Activity Kit	12
Hydrogen Peroxide (H ₂ O ₂) Colorimetric and Fluorescent Detection Kits	13
Superoxide Dismutase (SOD) Colorimetric Activity Kit	14
Thiol Fluorescent Detection Kit	15

ORDERING

- Online:** www.ArborAssays.com/order-form
- Phone:** Call 734-677-1774 or Toll Free: 855-677-1774. Monday-Friday 8:30 am to 5:30 pm, EST.
- Fax:** Send faxes to 734-677-6860.
- E-mail:** Send E-mail orders to Orders@ArborAssays.com
- Distributors:** Check our website at www.ArborAssays.com/distributors for a list of distributors.
- Mail:** Arbor Assays Inc., Sales Order Entry
1514 Eisenhower Place, Ann Arbor, MI 48108-3284, USA

Catalase Colorimetric & Fluorescent Activity Kits

Colorimetric Catalog No: K033-H1 (2 Plate)

Fluorescent Catalog No: K033-F1 (2 Plate)

FEATURES

- ▶ Use Measure Catalase Activity in Any Sample
- ▶ Time to Answer 45 Minutes
- ▶ Sensitivity Measure as Little as 0.052 U/mL
- ▶ Samples/Kit 89 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Format 96-well
- ▶ Species Species Independent
- ▶ Readout Colorimetric: 560 nm Fluorescent: 585 nm




SCIENTIFIC RELEVANCE

Hydrogen peroxide, (H₂O₂) is one of the most frequently occurring reactive oxygen species. It is formed either in the environment, as a by-product of aerobic metabolism, superoxide formation and dismutation, or as a product of oxidase activity. Both excessive hydrogen peroxide and its decomposition product hydroxyl radical, are harmful for most cell components. Its rapid removal is essential for all aerobically living prokaryotic and eukaryotic cells. One of the most efficient ways of removing peroxide is through the enzyme catalase, which is encoded by a single gene and is highly conserved among species. Mammals, including humans and mice, express catalase in all tissues. A high concentration of catalase can be found in the liver, kidneys and erythrocytes. The expression is regulated at transcription, post-transcription and post-translation levels. High catalase activity is detected in peroxisomes.

Colorimetric



Fluorescent



Ceruloplasmin (Cp) Colorimetric Activity Kit

Catalog No: K035-H1 (2 Plate)

Patented


FEATURES

- ▶ Use Wilson’s Disease Marker
- ▶ Sample Urine and Serum
- ▶ Validation Humans, Felids, Polar Bears, Giant Panda
- ▶ Species Multiple species
- ▶ Time to Answer 60 minutes
- ▶ Format 96-well
- ▶ Samples/Kit 88 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout Colorimetric, 560 nm



SCIENTIFIC RELEVANCE

Ceruloplasmin (Cp) is a multicopper oxidase enzyme involved in the safe handling of oxygen in some metabolic pathways of vertebrates. It was denoted ceruloplasmin, literally meaning ‘a blue substance from plasma’. Ceruloplasmin belongs to the family of multicopper oxidases which are among the few enzymes able to bind molecular oxygen to perform its complete reduction to water. Cp found in serum is expressed in the liver, but it is also expressed in the brain, lung, spleen and testis. Aceruloplasminaemia is an autosomal recessive disorder of iron metabolism characterized by the complete absence of ceruloplasmin. Ceruloplasmin is also associated with reproduction. Copper-deficient female rats seem to be protected against mortality. This protection has been suggested to be provided by estrogens, since estrogens alter the subcellular distribution of copper in the liver, leading to an increase in plasma copper levels and subsequent ceruloplasmin synthesis.



Corticosterone EIA & CLIA Kits

EIA Catalog No: K014-H1 (1 Plate) K014-H5 (5 Plate)
 CLIA Catalog No: K014-C1 (1 Plate) K014-C5 (5 Plate)


FEATURES

- ▶ Use Stress Marker in as Little as 1 µL Serum or Plasma
- ▶ Sample Serum, Plasma, Hair, Feathers, Urine, Fecal, Respiratory Vapor, and TCM
- ▶ Validation Mice, Rats, Humans, Monkeys, Birds, Cats, Ungulates
- ▶ Time to Answer 1.5 Hours (EIA) or 2 Hours (CLIA)
- ▶ Format 96-well, Break-Apart Strip
- ▶ Species Species Independent
- ▶ Samples/Kit 38/230 (EIA) or 39/231 (CLIA) in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout EIA: 450 nm CLIA: Glow Luminescent



SCIENTIFIC RELEVANCE

Corticosterone (Kendall’s Compound ‘B’) is a glucocorticoid secreted by the cortex of the adrenal gland. It is produced in response to stimulation of the adrenal cortex by ACTH and is the precursor of aldosterone. Corticosterone is a major indicator of stress and is the major stress steroid produced in non-human mammals. Studies involving corticosterone and levels of stress include impairment of long term memory retrieval, chronic corticosterone elevation due to dietary restrictions and in response to burn injuries. In addition to stress levels, corticosterone is believed to play a decisive role in sleep-wake patterns.



Cortisol EIA Kits

Catalog No: K003-H1 (1 Plate) K003-H5 (5 Plate) Strip Plates
 Catalog No: K003-H1W (1 Plate) K003-H5W (5 Plate) Whole Plates


FEATURES

- ▶ Use Stress Marker in as Little as 1 μ L Serum or Plasma
- ▶ Sample Serum, Plasma, Saliva, Hair, Urine, Fecal, and Respiratory Vapor
- ▶ Validation Humans, Primates, Ungulates, Monkeys
- ▶ Time to Answer 1.5 Hours
- ▶ Format 96-well, Break-Apart Strip or Whole Plates
- ▶ Species Species Independent
- ▶ Samples/Kit 39 or 231 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout Colorimetric, 450 nm



SCIENTIFIC RELEVANCE

Cortisol (hydrocortisone, Kendall’s Compound ‘F’) is the primary glucocorticoid produced and secreted by the adrenal cortex. It is often referred to as the “stress hormone” as it affects blood pressure, blood sugar levels, and other actions of stress adaptation. Immunologically, cortisol functions as an important anti-inflammatory and plays a role in hypersensitivity, immunosuppression, and disease resistance. In the metabolic aspect, cortisol promotes gluconeogenesis, liver glycogen deposition, and the reduction of glucose utilization. Production of cortisol follows an ACTH-dependent circadian rhythm, with a peak level in the morning and decreasing levels throughout the day. All but 4% of serum cortisol is bound to proteins including corticosteroid binding globulin and serum albumin. Abnormal cortisol levels are being evaluated for correlation with a variety of different conditions, such as prostate cancer, depression, schizophrenia, Cushing’s Syndrome, and Addison’s Disease.



Cortisone EIA & CLIA Kits

EIA Catalog No: K017-H1 (1 Plate) K017-H5 (5 Plate)
 CLIA Catalog No: K017-C1 (1 Plate) K017-C5 (5 Plate)


FEATURES

- ▶ Use Stress Marker in as Little as 1 µL Serum or Plasma
- ▶ Sample Serum, Plasma, Saliva, Hair, Urine, and Fecal
- ▶ Validation Mice, Rats, Humans, Monkeys, Birds, Felids, Ungulates
- ▶ Time to Answer 2 Hours (EIA) or 2.5 Hours (CLIA)
- ▶ Format 96-well, Break-Apart Strip
- ▶ Species Species Independent
- ▶ Samples/Kit 40 or 232 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout EIA: 450 nm CLIA: Glow Luminescent



SCIENTIFIC RELEVANCE

Cortisone (Kendall’s Compound ‘E’) was identified by extraction from bovine suprarenal gland tissue. Compound E was soon identified as cortisone. Cortisol and cortisone vary due to the activity of two 11β-hydroxysteroid dehydrogenases (11β-HSD). 11β-HSD1 is found primarily in the liver where it converts cortisone to cortisol, while 11β-HSD2 is found in tissues such as the kidney where cortisol receptor binding is required. 11β-HSD2 deactivates cortisol to cortisone, prohibiting receptor activation. This glucocorticoid “shuttle” helps to initiate and regulate the anti-inflammatory response. Monitoring the ratio of cortisone to cortisol has applications in diabetes, obesity, metabolic syndrome, osteoporosis, and chronic fatigue syndrome in addition to adrenal diseases.



FRAP[™] (Ferric Reducing Antioxidant Power) Detection Kit

Catalog No: K043-H1 (2 Plate)

Patent Protected


FEATURES

- ▶ Use Measure Ferric Reducing Anti-Oxidant Potential (FRAP)
- ▶ Sample Serum, Plasma, Urine, Food, Cosmetics, Additives
- ▶ Samples/Kit 89 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Time to Answer 30 Minutes
- ▶ Readout Colorimetric, 560 nm



SCIENTIFIC RELEVANCE

Potentially harmful reactive oxygen species (ROS) are produced as a consequence of normal aerobic metabolism. “Free Radicals” (FR) are usually removed or inactivated *in vivo* by a team of antioxidants. They are chemically stable atoms and molecules, which have one or more free electrons. Almost all biomolecules may be attacked by reactive free radicals. Free radicals are responsible for many pathological processes, or they can be generated as the result of the pathological stage and cause important secondary damage to biological systems and cells. Connections between free radicals and some serious diseases, including Parkinson’s and Alzheimer’s diseases, atherosclerosis, heart attacks, and chronic fatigue syndrome, have been demonstrated. However, short-term oxidative stress, the unbalance between the formation and scavenging of the reactive oxygen species, may be important in the prevention of aging due to triggering of the process known as mitohormesis. On average, 65 – 70% of the population is excessively impacted by oxidative stress caused by FRs.



Glutathione (GSH) Colorimetric Detection Kit

Catalog No: K006-H1 (4 Plate) K006-H1C-H/L (200 Cuvette)


FEATURES

- ▶ Use Measure GSH/GSSG to Determine Oxidative Stress
- ▶ Sample Cells, RBCs, Serum, Plasma, Urine, and Tissue
- ▶ Sensitivity To 32 pmol/Sample
- ▶ Format 96-well or Cuvette
- ▶ Species Species Independent
- ▶ Samples/Kit 89 (Total and GSSG) in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout Colorimetric, 405 nm

**MULTI
SPECIES**

SCIENTIFIC RELEVANCE

Glutathione (L- γ -glutamyl-L-cysteinylglycine; GSH) is the highest concentration non-protein thiol in mammalian cells and is present in concentrations of 0.5 - 10 mM. GSH is a tripeptide that contains an unusual peptide linkage between the amine group of cysteine and the carboxyl group of the glutamate side-chain. It is an antioxidant, preventing damage to important cellular components caused by reactive oxygen species such as free radicals and peroxides. Glutathione reduces disulfide bonds formed within cytoplasmic proteins to cysteines by serving as an electron donor. In the process, glutathione is converted to its oxidized form, glutathione disulfide (GSSG). Glutathione is found mostly in its reduced form, since the enzyme that reverts it from its oxidized form, glutathione reductase, is constitutive and inducible upon oxidative stress. The ratio of reduced glutathione to oxidized glutathione within cells is often used as a measure of cellular toxicity.



Glutathione (GSH) Fluorescent Detection Kits

96 Well: Catalog No: K006-F1 (1 Plate) K006-F5 (5 Plate)
384 Well: Catalog No: K006-F1D (2 Plate)


FEATURES

- ▶ Use Measure GSH/GSSG
- ▶ Sample Cells, RBC's, Serum, Plasma, Urine, and Tissues
- ▶ Convenient Measures Free and Total GSH in Same Sample Well
- ▶ Species Species Independent
- ▶ Sensitivity 45 nM Free GSH, 48 nM Total GSH
- ▶ Samples/Kit 96-well kits: 39 or 231 in Duplicate
384-well kit: 183 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout Fluorescent, 510 nm



SCIENTIFIC RELEVANCE

Glutathione (L-γ-glutamyl-L-cysteinylglycine; GSH) is the highest concentration non-protein thiol in mammalian cells and is present in concentrations of 0.5 - 10 mM. GSH is a tripeptide that contains an unusual peptide linkage between the amine group of cysteine and the carboxyl group of the glutamate side-chain. It is an antioxidant, preventing damage to important cellular components caused by reactive oxygen species such as free radicals and peroxides. Glutathione reduces disulfide bonds formed within cytoplasmic proteins to cysteines by serving as an electron donor. In the process, glutathione is converted to its oxidized form, glutathione disulfide (GSSG). Glutathione is found mostly in its reduced form, since the enzyme that reverts it from its oxidized form, glutathione reductase, is constitutive and inducible upon oxidative stress. The ratio of reduced glutathione to oxidized glutathione within cells is often used as a measure of cellular toxicity.



Glutathione Reductase Fluorescent Activity Kit

Catalog No: K009-F1 (1 Plate)


FEATURES

- ▶ Use Measure GR activity
- ▶ Sample RBCs, Serum, Plasma, and Cells
- ▶ Convenient 20 minute End Point or Kinetic Assay
- ▶ Sensitivity 9 $\mu\text{U/mL}$, World's Most Sensitive
- ▶ Species Species Independent
- ▶ Samples/Kit 41 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout Fluorescent, 510 nm



SCIENTIFIC RELEVANCE

Glutathione reductase (GR) plays an indirect but essential role in the prevention of oxidative damage within the cell by helping to maintain appropriate levels of intracellular glutathione (GSH). GSH, in conjunction with the enzyme glutathione peroxidase (GP), is the acting reductant responsible for minimizing harmful hydrogen peroxide. The regeneration of GSH is catalyzed by GR. GR is a ubiquitous 100-120 kDa dimeric flavoprotein that catalyzes the reduction of oxidized glutathione (GSSG) to reduced glutathione, using β -nicotinamide dinucleotide phosphate (NADPH) as the hydrogen donor. NADPH has been suggested to also act as an indirectly operating antioxidant, given its role in the recycling of GSSG to GSH and thus maintaining the antioxidative power of glutathione.



Glutathione S-Transferase Fluorescent Activity Kit

Catalog No: K008-F1 (1 Plate)


FEATURES

- ▶ Use Fluorescent Detection of GST Activity
- ▶ Sample Serum, Plasma, and Cell Lysates
- ▶ Samples/Kit 40 in Duplicate
- ▶ Convenient 30 Minute End Point or Kinetic Assay
- ▶ Sensitive < 100 μ U of GST Activity
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout Fluorescent, 460 nm



SCIENTIFIC RELEVANCE

The Glutathione S-Transferase (GST) family of isozymes function to detoxify and neutralize a wide variety of electrophilic molecules by mediating their conjugation with reduced glutathione. Human GSTs are encoded by several gene families, and expressed in almost all tissues. Given its pivotal role in ameliorating oxidative stress/damage, GST activity has been repeatedly investigated as a biomarker for arthritis, asthma, COPD, and multiple forms of cancer, as well as an environmental marker. Examination of GST isoforms and activity in human cancers, tumors and tumor cell lines has revealed the predominance of the acidic pi class. Furthermore, this activity is thought to substantially contribute to the innate or acquired resistance of specific neoplasms to anticancer therapy.



Hydrogen Peroxide Colorimetric & Fluorescent Detection Kits

Colorimetric Catalog No: K034-H1 (2 Plate)

Fluorescent Catalog No: K034-F1 (2 Plate)

FEATURES


- ▶ Use Measure H_2O_2 in Any Sample
- ▶ Time to Answer 15 Minutes
- ▶ Sensitivity To 2 pmol (65 pg) H_2O_2
- ▶ Samples/Kit Colorimetric: 89 in Duplicate Fluorescent: 88 in Duplicate
- ▶ Readout Colorimetric: 560 nm Fluorescent: 585 nm




SCIENTIFIC RELEVANCE

In biological systems, incomplete reduction of O_2 during respiration produces superoxide anion ($O_2^{\cdot-}$), which is spontaneously or enzymatically dismutated by superoxide dismutase to H_2O_2 . Many cells produce low levels of $O_2^{\cdot-}$ and H_2O_2 in response to a variety of extracellular stimuli, such as cytokines (TGF- β 1, TNF- α , and various interleukins), peptide growth factors (PDGF; EGF, VEGF, bFGF, and insulin), the agonists of heterotrimeric G protein-coupled receptors (GPCR) such as angiotensin II, thrombin, lysophosphatidic acid, sphingosine 1-phosphate, histamine, and bradykinin, or by shear stress. The addition of exogenous H_2O_2 , or the intracellular production in response to receptor stimulation, affects the function of various proteins including protein kinases, protein phosphatases, transcription factors, phospholipases, ion channels, and G proteins. In 1894, Fenton described the oxidation of tartaric acid by Fe^{2+} and H_2O_2 , in which H_2O_2 and O_2 participate in the production of singlet oxygen and peroxyxynitrite. The generation of these species may be concurrent with reactions involving iron, and under some circumstances they might be important contributors to H_2O_2 toxicity.

Colorimetric Standard Curve



Fluorescent Standard Curve



Superoxide Dismutase (SOD) Colorimetric Activity Kit

Catalog No: K028-H1 (2 Plate)


FEATURES

- ▶ Use Oxidative Stress Determination
- ▶ Sample Serum, Plasma, Urine, Cells and Tissue, and RBC
- ▶ Species Human and Other Mammalian Species
- ▶ Samples/Kit 89 in Duplicate
- ▶ Time to Answer 20 Minutes



SCIENTIFIC RELEVANCE

Short-lived and highly reactive oxygen species (ROS) such as $O_2^{\cdot -}$ (superoxide), $\cdot OH$ (hydroxyl radical), and H_2O_2 (hydrogen peroxide) are continuously generated *in vivo*. The cellular levels of ROS are controlled by antioxidant enzymes and small molecule antioxidants. The major antioxidant enzymes, superoxide dismutases (SODs), including copper-zinc superoxide dismutase (Cu/ZnSOD), manganese superoxide dismutase (MnSOD), and extracellular superoxide dismutase (EC-SOD). All play a critical roles in scavenging $O_2^{\cdot -}$. Decreased SOD activity results in elevated level of superoxide which in turn leads to decreased NO and increased peroxynitrite concentrations. The major intracellular SOD is a 32-kDa copper and zinc containing homodimer (Cu/Zn SOD). The mitochondrial SOD (MnSOD) is a manganese-containing 93-kDa homotetramer that is synthesized in the cytoplasm and translocated to the inner matrix of mitochondria. EC-SOD is the primary extracellular SOD enzyme and is highly expressed in many organs. Increased SOD activity levels are seen in Downs Syndrome, while decreased activity is seen in diabetes, Alzheimer’s disease, rheumatoid arthritis, Parkinson’s disease, uremic anemia, atherosclerosis, some cancers, and thyroid dysfunction.



Thiol Fluorescent Detection Kit

Catalog No: K005-F1 (1 Plate)


FEATURES

- ▶ Use Measure Thiol Content of Recombinant Proteins
- ▶ Adaptable Measure Protein Thiols (SH) in 6M GuHCl Buffers
- ▶ Sensitivity 4.62 nM
- ▶ Time to Answer 30 Minutes
- ▶ Samples/Kit 39 in Duplicate
- ▶ Stability Non-Toxic, Liquid 4°C Stable Reagents
- ▶ Readout Fluorescent, 510 nm



SCIENTIFIC RELEVANCE

Free thiols in biological systems have important roles. Oxidatively modified thiol groups of cysteine residues are known to modulate the activity of a growing number of proteins. One of the most pressing problems is to accurately determine the extent of modification of specific amino acids, such as cysteine residues, in a complex protein sample, especially in the presence of chaotropic agents such as guanidine hydrochloride. Typical methods such as using Ellman's reagent have limited sensitivity requiring large quantities of purified recombinant or native protein.





www.ArborAssays.com
(734) 677-1774
info@ArborAssays.com
©2014

180403

WEB 180403